

ANSWERS TO ASSIGNED EXERCISES 1-72

1. (d.)

$$\lambda = \frac{1}{2} B_0 xy$$

2.

$$\begin{aligned} 4\pi\epsilon_0 \vec{E} &= \frac{q}{r^2} \hat{r} \\ \rho &= q\delta^3(\vec{r}) \\ \vec{B} &= 0 \\ \vec{J} &= 0 \end{aligned}$$

3.

$$\begin{aligned} 4\pi\epsilon_0 V' &= \frac{q}{r} \\ \vec{A}' &= 0 \end{aligned}$$

5.

It is always possible to find a gauge in which $V' = 0$, by using $\lambda = \int_0^t V(t') dt'$. But, in general, a gauge cannot be found in which $\vec{A}' = 0$, because that would force $\vec{B} = 0$.

6.

- (b.) The constant of proportionality is $1/3!$.
(c.)

$$\det A = \frac{1}{4!} \epsilon_{ijkl} A_{im} A_{jn} A_{kp} A_{lq} \epsilon_{mnpq}$$

10.

- (a.) No (the interval is spacelike).
(b.) Yes.

$$|\vec{r}_A'' - \vec{r}_B''| = \sqrt{2}$$

- (c.) No (the interval is spacelike).
(d.) No (the interval is timelike).
(e.) Yes.

$$c|t_E'' - t_D''| = \sqrt{7}$$

11. Λ is symmetric, and its independent elements are

$$\begin{pmatrix} \gamma & -\gamma\beta n_1 & -\gamma\beta n_2 & 0 \\ \gamma n_1^2 + n_2^2 & (\gamma - 1)n_1 n_2 & 0 & 0 \\ & \gamma n_2^2 + n_1^2 & 0 & 0 \\ & & & 1 \end{pmatrix}$$

12. The answers are the same.

15.

$$\begin{aligned} \eta_{\max} &= 10.34 \\ \beta_{\max} &= 1 - (2.09 \times 10^{-9}) \\ x_{\max} &= 2.84 \times 10^{20} \text{ m} = 29,900 \text{ light yr} \\ \Delta t &= 1.89 \times 10^{12} \text{ sec} = 59,850 \text{ yr} \end{aligned}$$

16. (b.)

$$\text{fraction decaying} = 1 - \exp(-L/\gamma_0\beta_0 c\tau)$$

17.

(a.)

$$L = \gamma\beta c\tau = \frac{4}{3}c\tau$$

(b.)

$$\psi = \cos^{-1}(2\beta^2 - 1) = 73.74^\circ$$

19. (b.) The dimensions of $d\sigma$ are Joules⁻².

24.

(a.)

$$I = 2ne\beta_0 cA$$

(b.)

$$\frac{1}{\mu_0} \vec{B} = \hat{\phi} \frac{ne\beta_0 cA}{\pi s}$$

(c.)

$$n'_+ = n/\gamma_0$$

(d.)

$$n'_- = n\gamma_0(1 + \beta_0^2)$$

(e.)

$$c\epsilon_0 \vec{E}' = -\hat{s} \gamma_0 \beta_0 \frac{ne\beta_0 cA}{\pi s}$$

26.

(a.)

$$F_{\mu\nu} F^{\mu\nu} = -\frac{2}{c^2} (|\vec{E}|^2 - |c\vec{B}|^2)$$

(b.)

$$F^{\mu\nu}G_{\mu\nu} = -\frac{4}{c}\vec{E} \cdot \vec{B}$$

(c.)

$$\vec{E} \perp \vec{B} \quad \text{and} \quad |\vec{E}| > |c\vec{B}|$$

28.

(a.)

$$\sinh \eta = \frac{F_0}{mc} t$$

(b.)

$$t_1 = \frac{mc}{F_0}$$

29.(c.) x does not increase linearly with t .**30.**(c.) z does increase linearly with t .**32.**

$$\vec{E}(\vec{r}, t) = \hat{z} \frac{\mu_0 q_0 c^3 t}{2\pi[(ct)^2 - s^2]^{3/2}} \quad \text{for } t > s/c$$

$$= 0 \quad \text{otherwise}$$

$$c\vec{B}(\vec{r}, t) = -\hat{\phi} \frac{\mu_0 q_0 c^2 s}{2\pi[(ct)^2 - s^2]^{3/2}} \quad \text{for } t > s/c$$

$$= 0 \quad \text{otherwise}$$

35.

(b.)

$$\vec{S} \rightarrow \frac{q^2 c}{16\pi^2 \epsilon_0} \frac{\hat{z}}{b^2 z^2} \quad \text{as } \gamma \rightarrow \infty$$

(c.) No.

37.

(a.)

$$4\pi\epsilon_0 V(\vec{r}) \cong \frac{4ak}{\pi r}$$

(b.)

$$4\pi\epsilon_0 V(\vec{r}) \cong \frac{2a^2 k \cos \theta}{\pi r^2}$$

(c.)

$$4\pi\epsilon_0 V(\vec{r}) \cong -\frac{2a^3 k}{\pi^2 r^3} (3 \cos^2 \theta - 1)$$

38.

(a.)

$$\frac{\epsilon_0}{q} V(\vec{r}) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \frac{Y_{lm}(\theta, \phi) Y_{lm}^*(\theta', \phi') r'^l}{(2l+1)r^{l+1}}$$

(b.)

$$\frac{4\pi\epsilon_0}{q} V(\vec{r}) \cong \frac{1}{r}$$

$$+ \frac{r'}{r^2} (\cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi'))$$

$$+ \frac{r'^2}{r^3} \left(\left(\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right) \left(\frac{3}{2} \cos^2 \theta' - \frac{1}{2} \right) \right.$$

$$+ 3 \sin \theta \sin \theta' \cos \theta \cos \theta' \cos(\phi - \phi')$$

$$\left. + \frac{3}{4} \sin^2 \theta \sin^2 \theta' \cos(2(\phi - \phi')) \right)$$

39.Place charges $+1, -4, +6, -4, +1$ at $z = 2, 1, 0, -1$, and -2 , respectively.**42.**

$$f(\theta, \phi) \propto \cos^2 \theta \sin^2 \theta$$

43.(a.) q_{22} and q_{2-2} do not vanish and have equal relative weight.(b.) E-type (TM) radiation of types E_{22} and E_{2-2} are emitted with equal relative weight.

(c.)

$$f(\theta, \phi) \propto \sin^2 \theta (1 - \sin^2 \theta \cos^2 2\phi)$$

vanishes in six directions: $\theta = 0, \theta = \pi$, and at $(\theta = \frac{\pi}{2}; \phi = 0, \frac{\pi}{2}, \pi, \frac{3\pi}{2})$.**47.**

(a.)

$$\omega' = \frac{2\pi\beta c}{\Delta z/\gamma} \quad \left(\gamma \equiv \frac{1}{\sqrt{1-\beta^2}} \right)$$

(b.)

$$\omega = \frac{\omega'}{\gamma(1-\beta)}$$

(c.)

$$\begin{aligned}\lambda &= \Delta z \frac{1 - \beta}{\beta} \\ &= \frac{\Delta z}{\gamma^2 \beta (1 + \beta)} \\ &\rightarrow \frac{\Delta z}{2\gamma^2} \text{ as } \beta \rightarrow 1\end{aligned}$$

(d.)

$$\begin{aligned}E &= \gamma m_e c^2 \approx \sqrt{\frac{\Delta z}{2\lambda}} m_e c^2 \\ &= 1.616 \text{ GeV}\end{aligned}$$

51.

(b.) The state of polarization is RH (LH) elliptical if $\text{Im } \beta < 0$ ($\text{Im } \beta > 0$).

(c.)

$$\begin{aligned}\sqrt{1 + |\beta|^2} \vec{J}_1 &= \begin{pmatrix} 1 \\ \beta \end{pmatrix} \\ &= -\text{Im } \beta \begin{pmatrix} 1 \\ -i \end{pmatrix} + \begin{pmatrix} 1 + \text{Im } \beta \\ \text{Re } \beta \end{pmatrix} \\ &= \text{Im } \beta \begin{pmatrix} 1 \\ i \end{pmatrix} + \begin{pmatrix} 1 - \text{Im } \beta \\ \text{Re } \beta \end{pmatrix}\end{aligned}$$

No, this concern would not invalidate the answer to (b.).

52.

$$\begin{aligned}I_{A+B} &= I_A + I_B + \\ &+ \sqrt{I_A I_B} \frac{\text{Re}(\alpha \gamma^* + \beta \delta^*)}{\sqrt{(|\alpha|^2 + |\beta|^2)(|\gamma|^2 + |\delta|^2)}}\end{aligned}$$

53.

(a.)

$$J = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$$

(c.) No, the twisted nematic cell absorbs \hat{y} polarized light, whereas the rotator rotates \hat{y} polarized light into \hat{x} polarized light.

56.

(a.) No, Pedrotti's Jones matrix completely absorbs LH circularly polarized light.

(b.) Light passes through a device consisting of

a QWP with slow axis at 45° with respect to \hat{x} , followed by an \hat{x} polarizer, followed by a QWP with slow axis at -45° . The Jones matrix for this combination is half of that in part (a.).

59.

(a.)

$$\begin{aligned}\begin{pmatrix} \mathcal{S}_0 \\ \mathcal{S}_1 \\ \mathcal{S}_2 \\ \mathcal{S}_3 \end{pmatrix} &\equiv \mathcal{S}_0 = \mathcal{S}_p + \mathcal{S}_n \\ &= \begin{pmatrix} \sqrt{\mathcal{S}_1^2 + \mathcal{S}_2^2 + \mathcal{S}_3^2} \\ \mathcal{S}_1 \\ \mathcal{S}_2 \\ \mathcal{S}_3 \end{pmatrix} + \\ &+ \begin{pmatrix} \mathcal{S}_0 - \sqrt{\mathcal{S}_1^2 + \mathcal{S}_2^2 + \mathcal{S}_3^2} \\ 0 \\ 0 \\ 0 \end{pmatrix}\end{aligned}$$

61.

(c.) There is no relationship between V and \mathcal{V} .

62.

(a.) Referring to the hint, $\delta = \frac{\pi}{6}$. Then, approximating Betelgeuse as two point sources separated by $\Delta\theta_{\text{Betelgeuse}} = 0.047$ arc sec,

$$\text{slit sep} = \frac{2\delta}{\pi} \frac{\lambda}{\Delta\theta_{\text{Betelgeuse}}} \approx 0.878 \text{ m}$$

(b.)

$$l_c = \frac{\lambda_0^2}{\Delta\lambda} \approx 19.5 \lambda_0 \approx 11.7 \mu\text{m}$$

64.

(b.)

$$\frac{n_3}{n_1} = \sqrt{\frac{n_T}{n_0}}$$

65.

(a.) For $n_L = 1.38$, $n_H = 2.35$, and $N = 6$,

$$T = 1 - \left(\frac{\left(\frac{n_L}{n_H}\right)^{2N} - 1}{\left(\frac{n_L}{n_H}\right)^{2N} + 1} \right)^2 \approx 0.0064$$

(b.) $T = 1$.

68. $I'/I_0 = 1/4$ because (see hint) integral (b.) is half of integral (a.).

69.

(d.)

$$\frac{I_N(\psi_x, \psi_y)}{I_1(0, 0)} = N^2 \frac{\sin^2 \left(\frac{Nkd}{2} \sin \psi_x \right)}{\left(N \sin \left(\frac{kd}{2} \sin \psi_x \right) \right)^2} \times \\ \times \text{sinc}^2 \left(\frac{ka}{2} \sin \psi_x \right) \text{sinc}^2 \left(\frac{kb}{2} \sin \psi_y \right)$$

70.

(a.)

$$\mathcal{R}(\psi_x) = \left(\frac{\sin(2kd \sin \psi_x)}{4 \sin \left(\frac{kd}{2} \sin \psi_x \right)} \right)^2$$

(b.)

$$\mathcal{R}(\psi_x) = \frac{1}{4} \left(\cos \left(\frac{kd}{2} \sin \psi_x \right) + \cos \left(\frac{3kd}{2} \sin \psi_x \right) \right)^2$$

(c.)

$$\mathcal{R}(\psi_x) = \cos^2 \left(\frac{kd}{2} \sin \psi_x \right) \cos^2 (kd \sin \psi_x)$$

(d.) The three results are equivalent.

71.

This fuzzy thick slit is a convolution of a thick slit of width $\frac{a}{2}$ and a thick slit of width $\frac{3a}{2}$:

$$\mathcal{R}(\psi_x) = \text{sinc}^2 \left(\frac{ka}{4} \sin \psi_x \right) \text{sinc}^2 \left(\frac{3ka}{4} \sin \psi_x \right)$$

72.

(a.) At $\psi_x = 0$ the wave is pure x polarized.

(b.) The waves from the two slits are polarized orthogonally (RHCP *vs.* LHCP) and do not interfere:

$$\mathcal{R}(\psi_x) = \text{sinc}^2 \left(\frac{ka}{2} \sin \psi_x \right) \\ \psi_x(1^{\text{st}} \text{ min}) = \arcsin \frac{2\pi}{ka}$$